


This article was downloaded by: [INFLIBNET India Order]

On: 16 October 2009

Access details: Access Details: [subscription number 909277354]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Coordination Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title-content=t713455674>

### Synthesis, characterization, and structure of a new cobalt(II) Schiff-base complex with quinoxaline-2-carboxalidene-2-amino-5-methylphenol

Manju Sebastian <sup>a</sup>; Vasudevan Arun <sup>a</sup>; Ponminiesary P. Robinson <sup>a</sup>; Pally Leeju <sup>a</sup>; Digna Varghese <sup>a</sup>; Gopalakrishnan Varsha <sup>a</sup>; Karukapadath K. Mohammed Yusuff <sup>a</sup>

<sup>a</sup> Department of Applied Chemistry, Cochin University of Science and Technology, Cochin 682022, Kerala, India

First Published on: 15 October 2009

**To cite this Article** Sebastian, Manju, Arun, Vasudevan, Robinson, Ponminiesary P., Leeju, Pally, Varghese, Digna, Varsha, Gopalakrishnan and Mohammed Yusuff, Karukapadath K. (2009) 'Synthesis, characterization, and structure of a new cobalt(II) Schiff-base complex with quinoxaline-2-carboxalidene-2-amino-5-methylphenol', *Journal of Coordination Chemistry*, 99999:1,

**To link to this Article:** DOI: 10.1080/00958970903342117

**URL:** <http://dx.doi.org/10.1080/00958970903342117>

## PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Synthesis, characterization, and structure of a new cobalt(II) Schiff-base complex with quinoxaline-2-carboxalidine-2-amino-5-methylphenol

MANJU SEBASTIAN, VASUDEVAN ARUN,  
PONMINIESSARY P. ROBINSON, PALLY LEEJU, DIGNA VARGHESE,  
GOPALAKRISHNAN VARSHA and  
KARUKAPADATH K. MOHAMMED YUSUFF\*

Department of Applied Chemistry, Cochin University of Science and Technology,  
Cochin 682022, Kerala, India

(Received 24 April 2009; in final form 24 July 2009)

The mononuclear cobalt(II) complex  $[\text{CoL}_2] \cdot \text{H}_2\text{O}$  (where HL is quinoxaline-2-carboxalidine-2-amino-5-methylphenol) has been prepared and characterized by elemental analysis, conductivity measurement, IR, UV-Vis spectroscopy, TG-DTA, and X-ray structure determination. The crystallographic study shows that cobalt(II) is distorted octahedral with each tridentate NNO Schiff base in a cis arrangement. The crystal exhibits a 2-D polymeric structure parallel to [010] plane, formed by O–H...N and O–H...O intermolecular hydrogen bonds and  $\pi$ – $\pi$  stacking interactions, as a racemic mixture of optical enantiomers. The ligand is a Schiff base derived from quinoxaline-2-carboxaldehyde.

**Keywords:** Quinoxaline-2-carboxaldehyde; 2-Amino-5-methylphenol; Crystal structure; Schiff base; Cobalt(II)

### 1. Introduction

Transition metal complexes with rigid ligands forming frameworks are good candidates for functional materials. Properties such as electronic switching, magnetic ordering, conductivity, and catalysis [1–5] have prompted chemists to design 1-, 2-, and 3-D frameworks spaced by rigid ligands forming wide channels or cavities. Oxygen and nitrogen donor Schiff bases are of particular interest because of their ability to form transition metal complexes with unusual configurations, structural lability, and sensitivity to molecular environments [6–8]. They have proven effective in constructing supramolecular architectures such as coordination polymers and helical assemblies [9–11], employed for modeling active sites in biological systems. Their metal complexes find wide applications as antibacterial, antiviral, and antimalarial agents [12–14], homogeneous and heterogeneous catalysts [15, 16], and magnetic materials [17]. Cobalt Schiff-base complexes have received attention because of their important biological

\*Corresponding author. Email: yusuff@cusat.ac.in

applications [18] and there is continuing interest in the synthesis of new Schiff bases and their cobalt complexes.

A literature search reveals that the Schiff base derived from quinoxaline-2-carboxaldehyde and 2-amino-5-methylphenol has not been reported. The nitrogens in the quinoxaline unit can be an acceptor for hydrogen bonding [19] and may lead to polymeric structures. The electronic environment in the metal complexes of this Schiff base might be different from those derived from salicylaldehyde. In this article, we describe the synthesis and characterization of quinoxaline-2-carboxaldehyde-2-amino-5-methylphenol and its cobalt(II) complex. The crystal structure of the complex has been investigated by single crystal X-ray diffraction.

## 2. Experimental

### 2.1. Materials and physical measurements

Cobalt(II) acetate tetrahydrate (Merck) and 2-amino-5-methylphenol (Aldrich) were used as supplied. All other chemicals were of analytical grade purity and used without purification. Quinoxaline-2-carboxaldehyde was prepared using the reported procedure [20, 21]. Microanalyses of the compounds were done with an Elementar Vario EL III CHNS elemental analyzer; estimation of cobalt was carried out on a Thermo Electron Corporation, M series Atomic Absorption Spectrophotometer. The molar conductivity of the complex was measured using a Systronic conductivity bridge type 305 in dimethylformamide (DMF). FT-IR spectra were recorded in KBr pellets with a JASCO FTIR 4100 spectrophotometer from 4000 to 400  $\text{cm}^{-1}$ . The electronic spectrum of the ligand was recorded on a Thermo Electron Nicolet evolution 300 UV-Vis spectrophotometer. The  $^1\text{H}$  NMR spectra were recorded in  $\text{CDCl}_3$  on a Bruker 250 NMR spectrometer. TG-DTA analysis was carried out under nitrogen with a heating rate of  $10^\circ\text{C min}^{-1}$  using a Perkin Elmer Pyris Diamond TG/DTA analyzer.

### 2.2. Synthesis of quinoxaline-2-carboxaldehyde-2-amino-5-methylphenol

The monobasic tridentate Schiff base was prepared by mixing quinoxaline-2-carboxaldehyde (1.58 g, 0.01 mol) and 2-amino-5-methylphenol (1.23 g, 0.01 mol) in 1:1 molar ratio in methanol. The mixture was stirred at room temperature ( $27^\circ\text{C} \pm 1^\circ\text{C}$ ) for 1 h. The yellow flakes obtained were filtered, recrystallized from alcohol, and the purity was checked by TLC. Yield: 90%. Anal. Calcd for  $\text{C}_{16}\text{H}_{13}\text{N}_3\text{O}$  (263.29) (%): C, 72.46; H, 4.21; N, 15.64. Found (%): C, 72.99; H, 4.98; N, 15.96.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 298 K):  $\delta = 2.37$  (s, 3H, C-CH<sub>3</sub>), 7.80–8.20 (8H, Ar-H), 8.98 (s, 1H, azomethine proton), 9.70 (s, 1H, Ar-OH).

### 2.3. Synthesis of $[\text{Co}(\text{L})_2] \cdot \text{H}_2\text{O}$

Cobalt(II) acetate tetrahydrate (0.5 mmol, 1.25 g) was dissolved in a minimum of methanol and then added to 1:1 chloroform:methanol (100 mL) solution of HL (1.0 mmol, 2.63 g). During the addition, the color of the solution changed to dark violet.

Table 1. Crystal data summary for [Co(L)<sub>2</sub>]·H<sub>2</sub>O.

Empirical formula	C <sub>32</sub> H <sub>26</sub> CoN <sub>6</sub> O <sub>3</sub>
Formula weight	601.52
Crystal size (mm <sup>3</sup> )	0.40 × 0.36 × 0.02
Wavelength (Å)	0.71073
Crystal system	Monoclinic
Space group	<i>P</i> 2 <sub>1</sub> / <i>n</i>
Unit cell and dimensions (Å, °)	
<i>a</i>	13.7959(15)
<i>b</i>	13.8622(15)
<i>c</i>	14.5161(16)
$\beta$	94.684(2)
Volume (Å <sup>3</sup> ), <i>Z</i>	2766.8(5), 4
Calculated density (g cm <sup>-3</sup> )	1.444
Absorption coefficient (mm <sup>-1</sup> )	0.666
<i>F</i> (000)	1244
$\theta$ range for data collection (°)	1.96–26.03
Reflections collected	27217
Independent reflection	5445 [ <i>R</i> (int) = 0.0841]
Max. and min. transmission	0.987 and 0.766
Data/restraints/parameters	5416/2/391
Goodness-of-fit on <i>F</i> <sup>2</sup>	1.205
Final <i>R</i> indices [ <i>I</i> > 2 $\sigma$ ( <i>I</i> )]	<i>R</i> <sub>1</sub> = 0.0980, <i>wR</i> <sub>2</sub> = 0.1840
<i>R</i> indices (all data)	<i>R</i> <sub>1</sub> = 0.1289, <i>wR</i> <sub>2</sub> = 0.1975

$$R_1 = \sum(|F_o| - |F_c|) / \sum |F_o|, wR_2 = [\sum w(F_o^2 - F_c^2)^2 / (F_o^2)^2]^{0.5}.$$

It was then refluxed on a water bath for 3 h and left to cool at room temperature. Single crystals suitable for X-ray diffraction were obtained after 2 weeks by slow evaporation of the filtrate; the crystals obtained were thin. Yield: 85%. Anal. Calcd for C<sub>32</sub>H<sub>26</sub>CoN<sub>6</sub>O<sub>3</sub> (601.52) (%): C, 63.73; H, 4.23; N, 14.24; Co, 9.80. Found (%): C, 63.90; H, 4.36; N, 13.97; Co, 9.75. *A*<sub>M</sub> (DMF, Mho cm<sup>2</sup> mol<sup>-1</sup>): 3.9.

## 2.4. X-ray crystallography

X-ray crystal structure determination was performed with a Bruker SMART APEX CCD X-ray diffractometer using graphite monochromated Mo-K $\alpha$  radiation ( $\lambda = 0.71073$  Å,  $\varphi$  and  $\omega$  scans). The data were reduced using SAINTPLUS [22] and a multiscan absorption correction using SADABS (Siemens area detector absorption) was performed [23]. The structure was solved using SHELXS-97 and full-matrix least-squares refinement against *F*<sup>2</sup> was carried out using SHELXL-97 in anisotropic approximation for nonhydrogen atoms [24]. All hydrogens were assigned on the basis of geometrical considerations and were allowed to ride upon the respective carbons. A summary of crystallographic data and refinement parameters is given in table 1.

## 3. Results and discussion

### 3.1. Characterization of the ligand

The Schiff base was obtained by the condensation of quinoxaline-2-carboxaldehyde and 2-amino-5-methyl phenol. This yellow compound is soluble in common organic solvents.

The proton NMR spectrum of the Schiff base was recorded in  $\text{CDCl}_3$  solution using tetramethylsilane as internal standard. The signal at 9.70 ppm corresponds to phenolic –OH proton. The azomethine proton appears as a singlet at 8.98 ppm, and all eight aromatic protons appear as a multiplet at 7.80–8.20 ppm [25]. The methyl group attached to the phenolic ring is at 2.37 ppm as a singlet. The IR spectrum of the ligand exhibits a broad band in the range 3300–3500  $\text{cm}^{-1}$  due to phenolic (O–H) stretching. A strong band at 1262  $\text{cm}^{-1}$  is due to  $\nu\text{C–O}$  (phenolic). Weak bands in the region 2800–3060  $\text{cm}^{-1}$  are related to C–H stretches. The  $\nu(\text{C=N})$  of azomethine and that of quinoxaline appear as a group of bands in the region 1570–1630  $\text{cm}^{-1}$ . The peaks at 1626 and 1579  $\text{cm}^{-1}$  have been assigned to the azomethine –C=N stretch and quinoxaline (C=N) stretch, respectively [25, 26].

The electronic spectrum of the Schiff base was taken in methanol ( $10^{-4}\text{M}$ ). The wavelength maxima are observed at 46,511, 42,553, 34,013, and 25,974  $\text{cm}^{-1}$ . The strong bands at 46,511 and 42,553  $\text{cm}^{-1}$  are attributed to quinoxaline  $\pi\text{--}\pi^*$  and phenyl  $\pi\text{--}\pi^*$  transitions, respectively. The band at 25,974  $\text{cm}^{-1}$  is due to  $n\text{--}\pi^*$  transition of azomethine and ring (C=N) [27, 28]. The analytical data and spectral properties are consistent with the expected structure of the Schiff base.

### 3.2. Characterization of the complex

The cobalt complex was obtained as shining violet crystals, stable in air and soluble in polar organic solvents such as ethanol, methanol, DMF, and dimethylsulfoxide (DMSO). The molar conductivity of the complex in DMF confirms the nonelectrolytic nature [29]. Microanalytical data of the complex is in agreement with calculated values for  $[\text{Co}(\text{L})_2] \cdot \text{H}_2\text{O}$ .

**3.2.1. Infrared spectra.** FTIR spectra of the ligand and complex are given in Supplementary material. Comparison of the IR spectra of the complex with those of the uncomplexed ligand gives information regarding bonding sites of the ligand. Peaks in the region 3500–3300  $\text{cm}^{-1}$  for the complex are due to (O–H) stretching vibrations of lattice water and indicate hydrogen bonding [16]. The band observed at 935  $\text{cm}^{-1}$  for the complex is attributed to bending vibration of  $\text{H}_2\text{O}$  [30]. The complex displays a medium peak at 1667  $\text{cm}^{-1}$  due to azomethine (C=N) stretching. The azomethine  $\nu(\text{C=N})$  band in the complex is shifted by about 41  $\text{cm}^{-1}$  to higher energy compared to the free ligand. This increase in azomethine stretching frequency on coordination might be due to the extensive delocalization of the  $\pi$ -electrons in fully conjugated Schiff-base ligand [16]. The (C=N) stretch of quinoxaline ring in the complex appears as a strong band at 1575  $\text{cm}^{-1}$  for coordination of metal to ring nitrogen with a small shift for this band ( $\sim 5\text{cm}^{-1}$ ) on complexation, which might be due to involvement of the quinoxaline ring nitrogen in coordination. Such small shifts have been observed for benzimidazole complexes [31]. The C–O (phenolic) stretching band is observed at 1251  $\text{cm}^{-1}$  in the complex. These spectral assignments show that the Schiff base is coordinated to cobalt through azomethine nitrogen, phenolic oxygen, and quinoxaline nitrogen [32]. Conclusive evidence of the bonding is also shown by the presence of new bands in the spectrum of the complex at 567 and 468  $\text{cm}^{-1}$ , which can be attributed to (Co–O) and (Co–N) stretching vibrations, respectively [16].

**3.2.2. Electronic spectra.** Solution electronic spectra of the complexes were recorded in methanol in the UV-Vis region (200–900 nm) (Supplementary material). The band at  $25,974\text{ cm}^{-1}$  of the ligand shifts to  $27,027\text{ cm}^{-1}$  on complexation suggesting coordination of azomethine nitrogen with cobalt. The band at  $18,553\text{ cm}^{-1}$  is due to the  ${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$  [33, 34]. The intensity of this d–d band is considerably increased possibly due to the influence of nearby intense charge-transfer transitions ( $\log \epsilon = 4.6$ ). The other d–d transitions are masked by the strong charge transfer band, as observed in similar Schiff-base complexes [35, 36].

**3.2.3. Thermal analysis.** TG-DTA curves (Supplementary material) show the mass loss corresponding to one water molecule at 50–120°C, due to loss of lattice water [16]. The organic part of the complex decomposes into two steps, the first lies in the range 300–393°C with a DTG peak at 343°C. DTA shows an endotherm at 348°C corresponding to this step. The mass loss (39%) corresponds to loss of two 2-amino-5-methylphenol parts of the two Schiff bases. The next stage of decomposition starts at 412°C and is continuous. Since the TG curve of the complex showed continuous degradation at this stage, we were not able to isolate the intermediates. The steady mass loss observed at this stage may be due to the removal of the quinoxaline of the ligand. The decomposition was not complete even at 800°C.

**3.2.4. Description of the crystal structure.** The molecular structure of the complex was solved at 298 K. The complex crystallizes in monoclinic crystal system with the space group  $P2_1/n$  and the unit cell contains four molecules of  $C_{32}H_{26}CoN_6O_3$ . Figure 1 gives the ORTEP diagram with atomic labeling scheme. The important interatomic distances and angles are listed in table 2. Two of the monoanionic




Figure 1. ORTEP diagram of  $[Co(C_{16}H_{13}N_3O)_2] \cdot H_2O$  showing the atom labeling scheme with 50% probability ellipsoids.

Table 2. Selected bond lengths (Å) and angles (°) for [Co(L)<sub>2</sub>]·H<sub>2</sub>O.

Co(1)–O(1)	2.031(4)	Co(1)–N(2)	2.256(4)
Co(1)–O(2)	2.045(4)	Co(1)–N(5)	2.310(4)
Co(1)–N(4)	2.049(4)	N(1)–C(8)	1.288(6)
Co(1)–N(1)	2.052(4)	N(4)–C(24)	1.278(6)
O(1)–Co(1)–O(2)	100.55(18)	O(2)–Co(1)–N(5)	153.46(15)
O(1)–Co(1)–N(4)	96.58(15)	N(4)–Co(1)–N(5)	74.89(16)
O(2)–Co(1)–N(4)	79.73(15)	N(1)–Co(1)–N(5)	106.11(15)
O(1)–Co(1)–N(1)	79.98(16)	N(2)–Co(1)–N(5)	92.47(14)
O(2)–Co(1)–N(1)	99.71(15)	C(1)–O(1)–Co(1)	112.90(3)
N(4)–Co(1)–N(1)	176.38(17)	C(17)–O(2)–Co(1)	112.90(3)
O(1)–Co(1)–N(2)	155.61(15)	C(8)–N(1)–Co(1)	118.70(4)
O(2)–Co(1)–N(2)	87.80(15)	C(2)–N(1)–Co(1)	113.30(3)
N(4)–Co(1)–N(2)	107.50(16)	C(24)–N(4)–C(18)	126.30(4)
N(1)–Co(1)–N(2)	76.01(16)	C(24)–N(4)–Co(1)	120.30(4)
O(1)–Co(1)–N(5)	90.03(16)	C(18)–N(4)–Co(1)	113.40(3)

Table 3. Selected hydrogen bond interactions in the complex.

D–H...A	Symmetry	Distances (Å)		Angles (°) D–H...A
		D...A	H...A	
O(3)–H(31)...O(2)	1–x, 1–y, 1–z	2.895(9)	2.01(8)	169(6)
O(3)–H(32)...N(3)	1/2+x, 3/2–y, –1/2+z	3.035(10)	2.51(8)	124(6)
C(3)–H(3)...N(6)	1/2+x, 3/2–y, 1/2+z	3.467(7)	2.62	152
C(8)–H(8)...N(6)	1/2+x, 3/2–y, 1/2+z	3.388(7)	2.53(5)	157.1(12)
C(14)–H(14)...N(4)	1/2+x, 3/2–y, 1/2+z	3.426(7)	2.57	153
C(24)–H(24)...O(1)	–x, 1–y, 1–z	3.041(7)	2.35(3)	139(3)

tridentate Schiff-base ligands wrap the cobalt(II) center in the *cis* mode with coordination of two oxygens and four nitrogens (N<sub>4</sub>O<sub>2</sub>) in a distorted octahedral geometry. The molecule has no symmetry elements other than a C<sub>2</sub> symmetry axis and is expected to exhibit optical isomerism. However, both enantiomers coexist in the unit cell, making the space group centrosymmetric. The two phenolic hydroxyls are deprotonated on complexation. The azomethine (C=N) bond lengths (1.288(6) and 1.279(6) Å) are within the range of theoretical value (1.285 Å) [37]. The complex geometry can be explained by a rigid structure resulting in O(2)–Co(1)–N(5) and O(1)–Co(1)–N(2) bond angles that deviate significantly from linearity, i.e. 153.46(15) and 155.61(15)°. In the coordination sphere, the Co–O bond lengths are 2.031(4) and 2.045(4) Å; Co–N bond lengths are 2.049(4), 2.052(4), 2.256(4), and 2.310(4) Å. The Co–N (quinoxaline) bonds are elongated compared to Co–N (azomethine) bonds probably due to rigidity of the Schiff base. The benzene rings of the symmetry related quinoxalines are stacked *via* π–π interactions, having a centroid to centroid distance of 3.532 Å. Molecules are held together by O(3)–H(2)...N(3) and O(3)–H(31)...O(2) intermolecular hydrogen bonding and weak inter and intramolecular hydrogen bonding (table 3) forming a 2-D polymeric structure parallel to the [010] plane. Figure 2 illustrates the unit cell packing and polymeric structure of the molecules in the complex.




Figure 2. Packing diagram showing the 2-D polymeric chain along crystallographic *b*-axis (hydrogen atoms are omitted for clarity).

#### 4. Conclusion

Cobalt(II) complex of a new Schiff base has been synthesized and characterized. The crystallographic study shows that the cobalt(II) coordination sphere is distorted octahedrally with a *cis* arrangement of  $N_4O_2$  donor set of two tridentate NNO Schiff bases. The molecules exhibit a 2-D polymeric structure parallel to the [010] plane, formed by  $O-H\cdots N$  and  $O-H\cdots O$  intermolecular hydrogen bonds and  $\pi-\pi$  stacking. The crystal is a racemic mixture of two crystallographically independent enantiomers of the complex. To the best of our knowledge, this is the first report of the crystal structure of a mononuclear metal complex of a Schiff base derived from quinoxaline-2-carboxaldehyde.

#### Supplementary material

Crystallographic data for the structural analysis have been deposited with the Cambridge Crystallographic Data Centre, CCDC No. 729051. Copies of this information may be obtained free of charge from The Director, CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK (Fax: +44-1223-336033; E-mail: [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk) or [www: http://www.ccdc.cam.ac.uk](http://www.ccdc.cam.ac.uk)).

#### Acknowledgements

The X-ray data were collected on the diffractometer facilities at the University of Hyderabad provided by the Department of Science and Technology. The authors thank

Department of Science and Technology, India, for using the Sophisticated Analytical Instrumentation Facility (SAIF) at STIC, Cochin University of Science and Technology, Cochin for elemental analyses. M. Sebastian thanks Kerala State Council for Science, Technology and Environment, Kerala and D. Varghese thanks Council of Scientific and Industrial Research, India for research fellowships. We thank the referee for the kind help in fixing hydrogen atoms of the water molecule in the complex.

## References

- [1] T. Albrecht, A. Guckian, J. Ulstrup, J.G. Vos. *Nano Lett.*, **5**, 1451 (2005).
- [2] B.J. Coe, N.R.M. Curati. *Comment. Inorg. Chem.*, **25**, 147 (2004).
- [3] Y. Sato, S. Ohkoshi, K. Arai, M. Tozawa, K. Hashimoto. *J. Am. Chem. Soc.*, **125**, 14590 (2003).
- [4] A. Aumuller, P. Erk, G. Klebe, S. Hunig, J.U. Schutz, H.P. Werner. *Angew. Chem. Int. Ed.*, **25**, 740 (1986).
- [5] S. Liu, S. Dolder, P. Franz, A. Neels, H. Stoeckli-Evans, S. Decurtins. *Inorg. Chem.*, **42**, 4801 (2003).
- [6] S. Yamada. *Coord. Chem. Rev.*, **537**, 190 (1999).
- [7] A.D. Garnovskii, I.S. Vasilchenko, D.A. Garnovskii, B.I. Kharisov. *J. Coord. Chem.*, **62**, 151 (2009).
- [8] N. Raman, S.J. Raja, A. Sakthivel. *J. Coord. Chem.*, **62**, 691 (2009).
- [9] M.S. Ibrahim, S.E.H. Etaiw. *Synth. React. Inorg. Met. Org. Chem.*, **34**, 629 (2005).
- [10] A. Lalehzari, J. Desper, C.J. Levy. *Inorg. Chem.*, **47**, 1120 (2008).
- [11] P.A. Vigato, S. Tamburini. *Coord. Chem. Rev.*, **248**, 1717 (2004).
- [12] M.S. Refat, I.M. El-Deen, Z.M. Anwer, S. El-Ghol. *J. Coord. Chem.*, **62**, 1709 (2009).
- [13] P.H. Wang, G.J. Keck, E.J. Lien, M.M.C. Lai. *J. Med. Chem.*, **33**, 608 (1990).
- [14] J. Ziegler, T. Schuerle, L. Pasierb, C. Kelly, A. Elamin, K.A. Cole, D.W. Wright. *Inorg. Chem.*, **39**, 3731 (2000).
- [15] P.S. Chittilappilly, N. Sridevi, K.K.M. Yusuff. *J. Mol. Catal. A Chem.*, **286**, 92 (2008).
- [16] V. Arun, N. Sridevi, P.P. Robinson, S. Manju, K.K.M. Yusuff. *J. Mol. Catal. A Chem.*, **304**, 191 (2009).
- [17] X. He, C.Z. Lu, C.D. Wu. *J. Coord. Chem.*, **59**, 977 (2006).
- [18] W.M. Coleman, L.T. Taylor. *J. Am. Chem. Soc.*, **93**, 5446 (1971).
- [19] A. Kadzewski, M. Gdaniec. *Acta Cryst.*, **E62**, o3498 (2006).
- [20] H. Ohle, J.J. Kruff. *Ber. Chem. Ges.*, **77B**, 507 (1944).
- [21] C.L. Leese, H.N. Rydon. *J. Chem. Soc.*, 303 (1955).
- [22] SAINTPLUS, Bruker AXS Inc., Madison, USA (2003).
- [23] G.M. Sheldrick. *SADABS, Program for Empirical Absorption Correction*, University of Göttingen, Germany (1996).
- [24] G.M. Sheldrick. *SHELX-97, Programs for Crystal Structure Analysis*, University of Göttingen, Germany (1997).
- [25] V. Arun, P.P. Robinson, S. Manju, P. Leeju, G. Varsha, V. Digna, K.K.M. Yusuff. *Dyes Pigment.*, **82**, 268 (2009).
- [26] L.J. Bellamy. *The Infra-red Spectra of Complex Molecules*, Methuen, London (1956).
- [27] S.F. Dyke, A.J. Floyd, M. Sainsbury, R.S. Theobald. *Organic Spectroscopy – An Introduction*, Longman, New York (1978).
- [28] M.A. Neelakantan, S.S. Marriappan, J. Dharmaraja, T. Jeyakumar, K. Muthukumar. *Spectrochim. Acta A*, **71**, 628 (2008).
- [29] W.J. Geary. *Coord. Chem. Rev.*, **7**, 81 (1971).
- [30] K. Nakamoto. *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, 4th Edn, John Wiley and Sons Inc., New York (1986).
- [31] K.S. Bose, B.C. Sharma, C.C. Patel. *J. Inorg. Nucl. Chem.*, **32**, 1742 (1970).
- [32] E. Canpolat, M. Kaya. *J. Coord. Chem.*, **57**, 1217 (2004).
- [33] K.K.M. Yusuff, R. Sreekala. *Synth. React. Inorg. Met. Org. Chem.*, **21**, 553 (1991).
- [34] A.B.P. Lever. *Inorganic Electronic Spectroscopy*, 2nd Edn, Elsevier, Amsterdam (1984).
- [35] A.D. Naik, S.M. Annigeri, U.B. Gangadharmath, V.K. Revankar, V.B. Mahale. *Spectrochim. Acta A*, **58**, 1713 (2002).
- [36] S. Hong, X. Liang, H. Fang, X. Zhan, Z. Zhou, L. Chen, Y. Cai. *Transition Met. Chem.*, **34**, 115 (2009).
- [37] N.B.M. Elmkacher, M. Rzaigui, F. Bouachir. *Acta Cryst.*, **E64**, o271 (2008).